

Environmental geochemistry of Argemela Mine area (Barco, Central Portugal)

E.G. PEREIRA, I.M.H.R. ANTUNES

AND M.C.H.M. MONTEIRO

Polytechnic Institute of Castelo Branco, 6000-919 Castelo Branco, Portugal (tita@gmail.pt, imantunes@esa.ipcb.pt, carmoh@esa.ipcb.pt)

Mining and mineral-processing activities affect relatively small areas, but they can have a large local impact on the environment. The Argemela mining area is located at Central Iberian Zone, central Portugal. The mineralized veins intersect the Cambrian schist-metagraywacke Complex and is situated at about 13 km, to the east of the famous wolfram Portuguese producer, Minas da Panasqueira. Argemela and Panasqueira mines are included in the metalogenetic Sn-W province of Portugal. The Sn veins contain mainly quartz, albite, moscovite, lepidolite, cassiterite, columbite group minerals, amblygonite-montebrazite, pyrite, arsenopyrite and pyrrhotite. At Argemela mine, the underground exploration of cassiterite was dominant and ceased since fifty years ago. The tailings and rejected materials were deposited on the ground and are exposed to the air and water that can change the environmental geochemistry of the area.

The geochemical soil data show anomalies in the dependency of the mineralizations and mining activities.

Representative soil samples of the local geochemical background and soils collected inside mine influence were analysed. Soils inside mine influence present higher electrical conductivity (62 μ S/cm), Cd (28 mg/Kg), Pb (21 mg/Kg), Cu (4990 mg/Kg) and Zn (1168 mg/Kg) contents than soils collected outside mine influence (EC= 29 μ S/cm; Cd = 0.3 mg/Kg; Pb = 2 mg/Kg; Cu = 35 mg/Kg; Zn = 51 mg/Kg).

Most of soils are contaminated in Cu and Zn and must not be used for agriculture and human residences. Plants growing on the abandoned mine area have been study for their biogeochemical indication and mine remediation potential.

The obtained results indicate that *Erica cinerea* L., *Cytisus striatus* and *Cistus Ladanifer* are adapted to enriched Cu, Zn and Ni soils. *Erica cinerea* roofs tend to accumulate Cu (27 mg/Kg), while leaves from *Cytisus striatus* (9 mg/Kg) presents the highest Cu content of this species. Leaves from *Erica cinerea* L. and *Cytisus striatus* contain higher Zn, Cr, Ni and Pb contents than respectively roofs and stalks. The informations obtained for these species allows biogeochemical delineation of areas from similar soil anomalies.