

SEROPREVALENCE OF WEST NILE VIRUS ANTIBODIES IN WILD BIRDS OF PREY IN PORTUGAL

Filipa Loureiro^{1,2*}, Luís Cardoso^{2,3}, Ana C. Matos^{4,5}, Manuela Matos⁶ and Ana C. Coelho^{2,3}

¹Wildlife Rehabilitation Centre (CRAS), Veterinary Teaching Hospital (HVUTAD), University of Trás-os-Montes e Alto Douro (UTAD), Vila Real, PT; ²Animal and Veterinary Research Centre (CECAV), Associate Laboratory for Animal and Veterinary Sciences (AL4AnimalS), UTAD, Vila Real, PT; ³Department of Veterinary Sciences, School of Agrarian and Veterinary Sciences (ECAV), UTAD, Vila Real, PT; ⁴Research Centre for Natural Resources, Environment and Society (CERNAS), Polytechnic Institute of Castelo Branco, Castelo Branco, PT; ⁵Quality of Life in the Rural World (Q-RURAL), Polytechnic Institute of Castelo Branco, Castelo Branco, Portugal; ⁶Centre for the Research and Technology of Agro-Environmental and Biological Sciences (CITAB), UTAD, Vila Real, PT

This work was supported by the project UIDB/00772/2020 (doi:10.54499/UIDB/00772/2020) funded by the Portuguese Foundation for Science and Technology (FCT).

Introduction

West Nile virus (WNV) is one of the most widely distributed arboviruses in the world with zoonotic potential. Seroepidemiological studies are necessary to establish proper prevention and control strategies.

Materials and methods

WNV-total antibodies were searched in blood serum using an ELISA commercial kit. Samples from 136 individuals of diurnal raptors were analysed, from eight distinct Accipitriformes species. Eleven nocturnal birds of prey, belonging to four different species of Strigiformes were tested.

Discussion and conclusion

Despite the small nocturnal raptors sample, the seropositivity percentage was similar between the two groups. WNV is transmitted to raptors mainly through the bite of infected mosquitoes, although oral transmission has already been described.¹ Regarding nocturnal raptors, vectors are also the most likely transmission route, suggesting that louse flies, common avian hematophagous parasites, may be involved.² There is evidence that the mosquito population plays a key role in the transmission of zoonotic vector-borne pathogens.³ Studying how the mosquito community affects the seroprevalence of WNV in both groups would be an interesting follow-up.

In birds, as in mammals, protection against WNV is determined by the presence of antibodies in the blood. As antibodies are developed following exposure and infection with WNV in adult birds, detecting such antibodies may suggest disease, past subclinical infection, or recovery from a non-fatal infection. Furthermore, cross-reactivity between related flaviviruses may occur.^{1,4} A confirmation test, such viral neutralization test (VNT), would be recommended.


Results

In this study, 47 samples were positive (34.56%; 95% CI: 26.62-43.19%). A seropositivity of 31.62% (n=43; 95% CI: 23.92-40.14%) was found in diurnal, and 36.36% (n=4; 95%: 0.81-7.36%) in nocturnal raptors.

Diurnal raptors	Tested (n)	Positives (n)	Positives (%)
<i>Aegypius monachus</i>	36	10	27,78
<i>Aquila adalberti</i>	1	0	0
<i>Aquila chrysaetus</i>	1	0	0
<i>Buteo buteo</i>	1	0	0
<i>Gyps fulvus</i>	131	30	22,90
<i>Hieraaetus pennatus</i>	2	2	100
<i>Milvus migrans</i>	2	0	0
<i>Milvus milvus</i>	1	1	100
Total	136	43	31,62

Nocturnal raptors	Tested (n)	Positives (n)	Positives (%)
<i>Athene noctua</i>	1	1	100
<i>Bubo bubo</i>	4	1	25,00
<i>Strix aluco</i>	4	2	50,00
<i>Tyto alba</i>	2	0	0
Total	11	4	36,36

Fig. 1 – ELISA results of the tested birds.



References

- Vidaña B, Busquets N, Napp S, Pérez-Ramírez E, Jiménez-Clavero MÁ, Johnson N. The role of birds of prey in West Nile virus epidemiology. *Vaccines* (Basel). 2020; 8:550.
- Gancz AY, Barker IK, Lindsay R, Dibbernardo A, McKeever K, Hunter B. West Nile virus outbreak in North American owls, Ontario, 2002. *Emerg Infect Dis*. 2004; 10:2135-2142.
- Martínez-de la Puente J, Ferraguti M, Ruiz S, Roiz D, Llorente F, Pérez-Ramírez E, Jiménez-Clavero MÁ, Soriguer R, Figuerola J. Mosquito community influences West Nile virus seroprevalence in wild birds: implications for the risk of spillover into human populations. *Sci Rep*. 2018; 8:2599.
- Ferraguti M, LA Puente JM, Soriguer R, Llorente F, Jiménez-Clavero MÁ, Figuerola J. West Nile virus-neutralizing antibodies in wild birds from southern Spain. *Epidemiol Infect*. 2016; 144:1907-11.